

Response to: 'Presence of anti-phospholipid antibodies in COVID-19: a case series study' by Amezcua-Guerra *et al*

We thank Amezcua-Guerra *et al* for their interest in our study reporting on the course of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) disease 2019 (COVID-19) in a case series of patients with systemic lupus erythematosus under long-term treatment with hydroxychloroquine.^{1,2} Complementary to our work, Amezcua-Guerra *et al* address the issue of anti-phospholipid antibodies (anti-PL abs) during the course of COVID-19. Indeed, despite adequate thromboprophylaxis, COVID-19 is associated with a high rates of venous, as well as arterial, thromboembolic events, in particular in patients hospitalised in an intensive care unit.³⁻⁵ This state of hypercoagulation has been linked to an important systemic inflammatory response syndrome, with elevated serum levels of fibrinogen, factor VIII and D-dimers.^{6,7} Several reports, including the study by Amezcua-Guerra *et al*² have emphasised the high frequency of serum anti-PL abs and lupus anti-coagulant (LA) in a case series of patients with severe COVID-19, however with surprisingly heterogeneous results.

Amezcua-Guerra *et al* report a high frequency (57%) of both conventional (ie, those included in the antiphospholipid syndrome (APS) classification criteria) and non-conventional anti-PL abs in patients with severe and critical COVID-19, which appear to be associated with a hyperinflammatory state. An association with pulmonary thromboembolism has also been suggested although this concerned only 2 (17%) of the 12 patients who had at least one type of circulating anti-PL abs.² More recently, Zuo *et al*, measuring serum levels of eight different types of anti-PL abs in 172 patients hospitalised with COVID-19, detected anti-cardiolipin (anti-CL) IgM in 23%, anti-PS/PT IgG in 24% and anti-PS/PT IgM in 18% of the patients, with at least one type of anti-PL abs present in 52%.⁸ In contrast, Galeano-Valle *et al* reported a very low prevalence of conventional serum anti-PL abs among patients experiencing venous thromboembolism during the course of severe COVID-19.⁹ The results from their study were confirmed and extended by Borghi *et al* who also reported a low prevalence of anti- β_2 GPI IgG, IgA and IgM in patients with COVID-19 at a frequency of 15.6, 6.6 and 9.0%, respectively, as well as anti-CL IgG (5.7%) or IgM (6.6%), that was not associated with major thrombotic events.¹⁰ In the latter study, anti-PL abs were mainly directed against β_2 GPI, but they displayed an epitope specificity different from that of anti-PL abs present in APS.¹⁰ The explanation for the observed discrepancy between the rates of anti-PL abs reported in these studies might rely on the possibility that their generation is linked to the severity of COVID-19. In this respect, Bertin *et al* reported in a cohort of 56 patients with COVID-19 that such differences were found for anti-CL IgG whose presence were significantly associated with a severe form of the disease.¹¹ This observation was confirmed by Xiao *et al*, who showed that anti-PL abs, mostly anti- β_2 GPI and aCL IgA, were detected in 47.0% of critically ill patients, but not in patients with non-critical conditions.¹² Surprisingly, in the latter study, LA was detectable in only 2 of 66 critically ill patients. The presence of multiple anti-PL abs with a moderate serum titres of at least one type of anti-PL ab, was found to be statistically associated with a higher incidence of cerebral infarction.¹² Of note, anti-PL abs were mainly of the IgA isotype which suggest a cross-reactivity and/or breakdown of mucosal immune

tolerance induced by SARS-CoV-2 because of the pulmonary and intestinal mucosal tropism of this virus.

Meanwhile, many studies have shown a significant presence of LA in patients with severe COVID-19, mainly in critically ill conditions. In the study of Bowles *et al*, 31 patients (14%) were shown to be positive for an LA assay in a series of 216 patients with severe COVID-19 with only two patients having a confirmed or suspected venous thrombosis.⁷ Harzallah *et al* reported 25 patients (45%) positive for LA in a series of 56 critically ill patients with COVID-19,¹³ whereas in the study of Helm *et al*, 50 patients (33%) tested positive for an LA assay in a series of 150 patients with COVID-19-related acute respiratory disease syndrome (ARDS).³ These important rates of LA in critically ill patients with COVID-19 should however be interpreted with caution. Indeed, the extrapolation of LA results from patients receiving anticoagulants, which is now current clinical practice in the vast majority of patients hospitalised for COVID-19, is subject to discussion.¹⁴ Furthermore, one should be aware of false-positive LA testing results in patients with COVID-19 because many assays are sensitive to the presence of C-reactive-protein resulting in false positives.¹⁵

We recently reported, on a series of 25 patients with refractory COVID-19-related ARDS, 23 cases (92%) of LA. Anti-CL or anti- β_2 GPI abs were observed in 13 (52%) and 3 (12%) cases, respectively.¹⁶ Three patients (12%) were triple positive for LA, anti-CL and anti- β_2 GPI abs, whereas massive pulmonary embolism was diagnosed in six patients, all positive for the presence of anti-PL abs.

During acute infection, thrombosis or inflammation, serum levels of different anti-PL abs may transiently arise. Strikingly, however, this elevation of anti-PL abs and/or LA titres reported in a major proportion of patients with severe COVID-19 has rarely been observed in other pathologies. Nevertheless, the involvement of anti-PL abs in the induction of a hypercoagulable state and the possibility that SARS-Cov-2 may trigger the development of 'COVID-19-induced APS-like syndrome' have to be confirmed in large clinical series. Notwithstanding, the high frequency and wide variety of anti-PL abs observed in patients with COVID-19 cannot be ignored.

Alexis Mathian ¹, Marc Pineton De Chambrun ², Alain Combes,² Zahir Amoura¹

¹Sorbonne Université, Assistance Publique-Hôpitaux de Paris, Groupement Hospitalier Pitié-Salpêtrière, French National Referral Center for Systemic Lupus Erythematosus, Antiphospholipid Antibody Syndrome and Other Autoimmune Disorders, Service de Médecine Interne 2, Institut E3M, Inserm UMR5, Centre d'Immunologie et des Maladies Infectieuses (CIMI-Paris), Paris, France

²Sorbonne Université, Assistance Publique-Hôpitaux de Paris, Groupement Hospitalier Pitié-Salpêtrière, Institut de Cardiométabolisme et Nutrition (ICAN), Service de Médecine Intensive-Réanimation, Inserm UMR51166, Paris, France

Correspondence to Dr Alexis Mathian, Internal Medicine, University Hospital Pitié Salpêtrière, Paris, France; alexis.mathian@aphp.fr

Handling editor Josef S Smolen

Contributors AM, MPDC, AC and ZA wrote the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Commissioned; internally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful,

non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Mathian A, Pineton De Chambrun M, Combes A, *et al.*
Ann Rheum Dis Epub ahead of print: [please include Day Month Year]. doi:10.1136/annrheumdis-2020-218145

Received 21 July 2020
Accepted 22 July 2020



► <http://dx.doi.org/10.1136/annrheumdis-2020-218100>

Ann Rheum Dis 2020;**0**:1–2. doi:10.1136/annrheumdis-2020-218145

ORCID iDs

Alexis Mathian <http://orcid.org/0000-0002-7653-6528>

Marc Pineton De Chambrun <http://orcid.org/0000-0002-6321-858X>

REFERENCES

- Mathian A, Mahevas M, Rohmer J, *et al.* Clinical course of coronavirus disease 2019 (COVID-19) in a series of 17 patients with systemic lupus erythematosus under long-term treatment with hydroxychloroquine. *Ann Rheum Dis* 2020. 10.1136/annrheumdis-2020-217847
- Amezcu-Guerra LM, Rojas-Velasco G, Brianza-Padilla M, *et al.* Presence of antiphospholipid antibodies in COVID-19: a case series study. *Ann Rheum Dis* 2020. 10.1136/annrheumdis-2020-218100
- Helms J, Tacquard C, Severac F, *et al.* High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study. *Intensive Care Med* 2020;46:1089–98.
- Klok FA, Kruij MJHA, van der Meer NJM, *et al.* Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res* 2020;191:145–7.
- Llitjos J-F, Leclerc M, Chochois C, *et al.* High incidence of venous thromboembolic events in anticoagulated severe COVID-19 patients. *J Thromb Haemost* 2020;18:1743–6.
- Wang D, Hu B, Hu C, *et al.* Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020. doi:10.1001/jama.2020.1585. [Epub ahead of print: 07 Feb 2020].
- Bowles L, Platton S, Yartey N, *et al.* Lupus anticoagulant and abnormal coagulation tests in patients with Covid-19. *N Engl J Med* 2020;383:288–90.
- Zuo Y, Estes SK, Gandhi AA, *et al.* Prothrombotic antiphospholipid antibodies in COVID-19. *medRxiv*2020.
- Galeano-Valle F, Oblitas CM, Ferreiro-Mazón MM, *et al.* Antiphospholipid antibodies are not elevated in patients with severe COVID-19 pneumonia and venous thromboembolism. *Thromb Res* 2020;192:113–5.
- Borghini MO, Beltagy A, Garrafa E, *et al.* Prevalence, specificity, and clinical association of anti-phospholipid antibodies in COVID-19 patients: are the antibodies really guilty? *medRxiv* 2020. doi:10.1101/2020.06.17.20134114. [Epub ahead of print: 19 Jun 2020].
- Bertin D, Brodovitch A, Beziane A, *et al.* Anti-Cardiolipin IgG autoantibodies are an independent risk factor of COVID-19 severity. *Arthritis Rheumatol* 2020. doi:10.1002/art.41409. [Epub ahead of print: 21 Jun 2020].
- Xiao M, Zhang Y, Zhang S, *et al.* Brief report: Anti-phospholipid antibodies in critically ill patients with coronavirus disease 2019 (COVID-19). *Arthritis Rheumatol* 2020.
- Harzallah I, Debliquis A, Drénou B. Lupus anticoagulant is frequent in patients with Covid-19. *J Thromb Haemost* 2020. doi:10.1111/jth.14867. [Epub ahead of print: 23 Apr 2020].
- Pengo V, Tripodi A, Reber G, *et al.* Update of the guidelines for lupus anticoagulant detection. Subcommittee on Lupus Anticoagulant/antiphospholipid Antibody of the Scientific and Standardisation Committee of the International Society on Thrombosis and Haemostasis. *J Thromb Haemost* 2009;7:1737–40.
- Schouwens SME, Delanghe JR, Devreese KJM. Lupus anticoagulant (lac) testing in patients with inflammatory status: does C-reactive protein interfere with lac test results? *Thromb Res* 2010;125:102–4.
- Pineton de Chambrun M, Frere C, Miyara M, *et al.* High frequency of antiphospholipid antibodies in critically ill COVID-19 patients: a link with hypercoagulability? *J Intern Med* 2020. doi:10.1111/joim.13126. [Epub ahead of print: 12 Jun 2020].